

**MASTER**

Los Alamos National Laboratory is operated by the University of California for the United States Department of Energy under contract W-7405-ENG-36

TITLE MEASURING THE ANTIQUARK DISTRIBUTION FUNCTION IN NUCLEI

LA-UR--86-2686

DE86 015350

AUTHOR(S) Gerald T(homas) GARVEY

SUBMITTED TO International Nuclear Physics Conference  
Harrogate, United Kingdom  
August 25-30, 1986  
Presentation to and inclusion in meeting proceedings.

**DISCLAIMER**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

By acceptance of this article, the publisher recognizes that the U.S. Government retains a nonexclusive, royalty free license to publish or reproduce the published form of this contribution or to allow others to do so for U.S. Government purposes.

The Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy.

**Los Alamos** Los Alamos National Laboratory  
Los Alamos, New Mexico 87545

jsw

## MEASURING THE ANTIQUARK DISTRIBUTION FUNCTION IN NUCLEI

G. T. Garvey

Los Alamos National Laboratory, Los Alamos, NM 87545, U.S.A.

The EMC<sup>1</sup> effect has prompted a great deal of speculation on possible nuclear medium effects on hadron structure. This speculation has continued unabated and unresolved from the first publication of the result. Part of the difficulty arises from the limited information available from inclusive deep-inelastic charged lepton scattering. The structure function extracted from such experiments is

$$F_2^{em}(x) = \sum_f e_f^2 (q_f(x) + \bar{q}_f(x)) \quad (1)$$

where  $f$  is the quark flavor ( $u, d, \dots$ ) and  $q_f(x)$  is the probability of a quark of type  $f$  carrying a fraction  $x$  of the total momentum of a nucleus. As such it is insensitive to quark type, coupling only via the charge squared.

As inclusive neutrino reactions<sup>2</sup> are not able to obtain sufficient statistics to provide additional specific information on quark momentum densities several authors<sup>3</sup> have advocated using the Drell-Yan process to supplement our existing knowledge. The Drell-Yan yield determines the following quantity

$$F(x_1, x_2) = \sum_f e_f^2 [q_f(x_1)\bar{q}_f(x_2) + \bar{q}_f(x_1)q_f(x_2)] \quad (2)$$

where  $x_1$  refers to the incident hadron and  $x_2$  to a target hadron. Using information on the structure function of incident hadron ( $p, \pi, k$ , etc.) one can select specific aspects of the target quark distribution. For example using incident protons and selecting large  $x_1$  ( $x_1 > 0.6$ ) an antiquark distribution function of the target can be measured. Recent theoretical and experimental developments indicate that the Drell-Yan process can be reliably used to measure ratios of quark distribution functions in different nuclei. An approved experiment<sup>4</sup> at FNAL to measure the ratio of antiquark distribution functions in Fe and D at low  $x_2$  ( $0.05 < x_2 < 0.3$ ) will be briefly discussed. Arguments will be presented that show proton energies lower than 45 GeV are not useful in Drell-Yan studies of nuclear effects.

### References

1. J. J. Aubert et al., Phys. Lett. 123B (1983) 275.
2. H. Abramowicz et al., Zeit. für Phys. C25 (1984) 29.
3. e.g., R. P. Bickerstaff et al., Phys. Rev. D33 (1986) 3228; E. Berger, Nucl. Phys. B267 (1986) 231.
4. FNAL E-772, Collaboration of Fermi National Accelerator Laboratory, Los Alamos National Laboratory, University of Illinois (Chicago), State University of New York at Stony Brook, University of Texas, and University of Washington.